

Poster presentation

Open Access

## European mania in bipolar longitudinal evaluation of medication (EMBLEM) study: Greek baseline results of a two year European, observational health outcomes study in bipolar disorder

Charalambos Touloumis<sup>\*1</sup>, Vangelis Drossinos<sup>2</sup>, Maria Kyrana<sup>2</sup> and Dimitrios Massaras<sup>2</sup>

Address: <sup>1</sup>Psychiatric Hospital of Attica "Dafni", Greece and <sup>2</sup>Eli Lilly and Company, Greece

\* Corresponding author

from International Society on Brain and Behaviour: 2nd International Congress on Brain and Behaviour  
Thessaloniki, Greece. 17–20 November 2005

Published: 28 February 2006

*Annals of General Psychiatry* 2006, **5**(Suppl 1):S198 doi:10.1186/1744-859X-5-S1-S198

### Background

Although several studies have examined the treatment of mania in routine clinical settings in individual countries, there has yet to be an observational study of actual care to evaluate the short and long-term management of patients with mania across Europe in terms of clinical, functional and economic outcomes.

EMBLEM (European Mania in Bipolar Evaluation of Medication) is a 2-year prospective observational study on the outcomes of pharmacological treatment for acute mania in the context of bipolar disorder. It has been successfully implemented in 14 European countries since its start in December 2002. Enrollment into the study ended on June 30, 2004. Follow-up observations will continue until July 2006.

The current objective is to describe the baseline treatment patterns of 634 patients enrolled into EMBLEM during an acute manic episode.

### Materials and methods

EMBLEM is a 2-year prospective, Pan-European observational study on the outcomes of pharmacological treatment for mania in the context of bipolar disorder. Adult patients are enrolled if they: present within the standard course of care as in- or outpatient; have, at the discretion of the treating psychiatrist, initiated or changed oral medication (antipsychotic, lithium or anticonvulsant) for treatment of acute mania; are not participating in an interventional study.

Investigators were asked, but not required, to include 50% of patients into the study who were initiated or changed to olanzapine and 50% of patients initiated or changed to non-olanzapine treatment. 58 psychiatrists enrolled 663 patients in Greece using the same study methods including assessment of mood with CGI-BP mania and CGI-BP depression rating scales as well as the Life Chart Method.

### Results

Baseline analysis has been completed with the last patient having been enrolled in June 2004. We will present baseline data of 634 patients including suicide history, treatment patterns prior to and after start of new medication as well as the reason for start of new oral medication.

### Discussion

The 663 patients from Greece represent the 18% of the total number of patients having been enrolled in EMBLEM.

1/4 of the patients had at least once attempted suicide.

1/4 of the patients were not taking any anti-manic medication when entering into the study.

Lack of effectiveness is the most common reason for switching oral medication.

Atypicals represented the 86% of the medication (antipsychotics, lithium or anticonvulsants) as monotherapy following the start of new oral medication.

## References

1. Haro JM, et al.: **European mania in bipolar longitudinal evaluation of medication (EMBLEM) study: study design and recruitment of a 2-year, pan-European, observational health outcomes study in bipolar disorder.** *Bipolar Disord* 2003, **5**(Suppl 1):P82.
2. Wells KB: **Treatment research at the crossroads; the scientific interface of clinical trials and effectiveness research.** *Am J Psychiatry* 1999, **156**:5-10.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

